LIBRARY
OF THE
UNIVERSITY
OF ILLINOIS
590.5
FI
v.34
MAY 5 1964
BIOLOGY.

Return this book on or before the
Latest Date stamped below.

Theft, mutilation, and underlining of books are reasons for disciplinary action and may result in dismissal from the University.

University of Illinois Library

JAN 12 1966

DEC 4 1967

SEP 15 1971

NOV 15 1976

DEC 14 1990

DEC 07 1992

L161—O-1096

590.5
F1
no. 34 32

FIELDIANA · ZOOLOGY

Published by
CHICAGO NATURAL HISTORY MUSEUM

Volume 34 | NOVEMBER 10, 1955 | No. 32

A NEW SPECIES OF CHONDRIA WITH A KEY TO THE BORNEAN SPECIES (COLEOPTERA: ENDOMYCHIDAE)

H. F. STROHECKER
UNIVERSITY OF MIAMI, CORAL GABLES, FLORIDA

The following new species of endomychid beetle was collected by the Bornean Zoological Expedition (1950) of Chicago Natural History Museum.

Chondria cardiaca sp. nov. Figure 64.

Broadly cordiform, ferruginous except for the eyes and joints 9 and 10 of the antennae, which are black. Length 2.4–2.7 mm. In the male the antennae are about half as long as the body, with joints 3–8 each longer than broad, subequal in length but increasing in diameter distally, joint 9 about as long as broad, 10 transverse, 11 oval and about equal in length to 9 and 10 together. Pronotum transverse, narrowed from base to apex with only moderate curvature of the sides, its front angles somewhat produced and acute, its raised margins narrowed behind, the hind margin deeply notched internal to the hind angles. The basal sulcus of the pronotum is not deep but very evident and ends on each side in a punctiform fovea. Elytra strongly convex, cordiform, the umbones prominent. Hind femur with a flange that is produced distally into a slender, acute tooth.

The female is similar to the male but antennal joints 4–8 are no longer than broad, the front angles of the pronotum are right angles, the elytra are of shorter form and the hind femur lacks the flange and tooth found in the male.

Holotype.—Male collected by Robert F. Inger in the Sapagaya Forest Preserve, Sandakan District, British North Borneo, July 27, 1950. In Chicago Natural History Museum.

NAT. HIST.

THE LIBRARY OF THE
NOV 15 1955
UNIVERSITY OF ILLINOIS

Allotype.—Female collected by D. Dwight Davis at southeast end of Dewhurst Bay, Kinabatangan District, British North Borneo, July 21, 1950. In Chicago Natural History Museum.



FIG. 64. *Chondria cardiaca* sp. nov. Left, dorsal view, left half of female; above, hind femur of male.

Paratypes.—Two males and one female with the same data as the holotype, and four females with the same data as the allotype.

Three other species of *Chondria* from Borneo were described by Arrow in 1920 (Trans. Ent. Soc. London, pp. 58–61). The following is a key to the four species:

1. Elytra plainly seriately punctured 2
   Elytra not seriately punctured, cordiform ..... *cardiaca*
2. Club of antenna much shorter than stalk ..... *seriosetosa*
   Antennal club equal to or longer than stalk 3
3. Sides of pronotum serrulate ..... *globulosa*
   Sides of pronotum smooth; raised margin narrowed behind ..... *nitida*

90.5
I
34 32

UNIVERSITY OF ILLINOIS-URBANA
590.5FI C001
FIELDIANA, ZOOLOGY$CHGO
34 1952-57
3 0112 009379659